

# Propositional and Predicate Logic - X

Petr Gregor

KTIML MFF UK

WS 2023/2024

# Properties of theories

We introduce syntactic variants of previous semantical definitions.

Let  $T$  be a theory of a language  $L$ . If a sentence  $\varphi$  is provable from  $T$ , we say that  $\varphi$  is a *theorem* of  $T$ . The set of theorems of  $T$  is denoted by

$$\text{Thm}^L(T) = \{\varphi \in \text{Fm}_L \mid T \vdash \varphi\}.$$

We say that a theory  $T$  is

- *inconsistent* if  $T \vdash \perp$ , otherwise  $T$  is *consistent*,
- *complete* if it is consistent and every sentence is provable or refutable from  $T$ , i.e.  $T \vdash \varphi$  or  $T \vdash \neg\varphi$ .
- an *extension* of a theory  $T'$  of  $L'$  if  $L' \subseteq L$  and  $\text{Thm}^{L'}(T') \subseteq \text{Thm}^L(T)$ , we say that an extension  $T$  of a theory  $T'$  is *simple* if  $L = L'$ ; and *conservative* if  $\text{Thm}^{L'}(T') = \text{Thm}^L(T) \cap \text{Fm}_{L'}$ ,
- *equivalent* with a theory  $T'$  if  $T$  is an extension of  $T'$  and vice-versa.

# Corollaries

From the soundness and completeness of the tableau method it follows that these syntactic definitions agree with their semantic variants.

**Corollary** For every theory  $T$  and sentences  $\varphi, \psi$  of a language  $L$ ,

- $T \vdash \varphi$  if and only if  $T \models \varphi$ ,
- $\text{Thm}^L(T) = \theta^L(T)$ ,
- $T$  is inconsistent if and only if  $T$  is unsatisfiable, i.e. it has no model,
- $T$  is complete if and only if  $T$  is semantically complete, i.e. it has a single model, up to elementary equivalence,
- $T, \varphi \vdash \psi$  if and only if  $T \vdash \varphi \rightarrow \psi$  (*Deduction theorem*).

**Remark** Deduction theorem can be proved directly by transformations of tableaux.

## Existence of a countable model and compactness

**Theorem** Every consistent theory  $T$  of a countable language  $L$  without equality has a *countably infinite* model.

*Proof* Let  $\tau$  be the systematic tableau from  $T$  with  $F\perp$  in the root. Since  $\tau$  is finished and contains a noncontradictory branch  $V$  as  $\perp$  is not provable from  $T$ , there exists a *canonical model*  $\mathcal{A}$  from  $V$ . Since  $\mathcal{A}$  agrees with  $V$ , its reduct to the language  $L$  is a desired countably infinite model of  $T$ .  $\square$

*Remark* This is a weak version of so called *Löwenheim-Skolem theorem*. In a countable language with *equality* the canonical model with equality is *countable* (i.e. finite or countably infinite).

**Theorem** A theory  $T$  has a model iff every *finite* subset of  $T$  has a model.

*Proof* The implication from left to right is obvious. If  $T$  has no model, then it is inconsistent, i.e.  $\perp$  is provable by a systematic tableau  $\tau$  from  $T$ . Since  $\tau$  is finite,  $\perp$  is provable from some finite  $T' \subseteq T$ , i.e.  $T'$  has no model.  $\square$

## Non-standard model of natural numbers

Let  $\underline{\mathbb{N}} = \langle \mathbb{N}, S, +, \cdot, 0, \leq \rangle$  be the standard model of natural numbers.

Let  $\text{Th}(\underline{\mathbb{N}})$  denote the set of all **sentences** that are valid in  $\underline{\mathbb{N}}$ . For  $n \in \mathbb{N}$  let  $\underline{n}$  denote the term  $S(S(\dots(S(0))\dots))$ , so called the  *$n$ -th numeral*, where  $S$  is applied  $n$ -times.

Consider the following theory  $T$  where  $c$  is a new constant symbol.

$$T = \text{Th}(\underline{\mathbb{N}}) \cup \{ \underline{n} < c \mid n \in \mathbb{N} \}$$

*Observation* Every finite subset of  $T$  has a model.

Thus by the compactness theorem,  $T$  has a model  $\mathcal{A}$ . It is a *non-standard model of natural numbers*. Every sentence from  $\text{Th}(\underline{\mathbb{N}})$  is valid in  $\mathcal{A}$  but it contains an element  $c^{\mathcal{A}}$  that is greater than every  $n \in \mathbb{N}$  (i.e. the value of the term  $\underline{n}$  in  $\mathcal{A}$ ).

# Equisatisfiability

We will see that the problem of satisfiability can be *reduced* to open theories.

- Theories  $T$ ,  $T'$  are *equisatisfiable* if  $T$  has a model  $\Leftrightarrow T'$  has a model.
- A formula  $\varphi$  is in the *prenex (normal) form (PNF)* if it is written as

$$(Q_1x_1) \dots (Q_nx_n)\varphi',$$

where  $Q_i$  denotes  $\forall$  or  $\exists$ , variables  $x_1, \dots, x_n$  are all distinct and  $\varphi'$  is an open formula, called the *matrix*.  $(Q_1x_1) \dots (Q_nx_n)$  is called the *prefix*.

- In particular, if all quantifiers are  $\forall$ , then  $\varphi$  is a *universal* formula.

To find an open theory equisatisfiable with  $T$  we proceed as follows.

- We replace axioms of  $T$  by equivalent formulas in the *prenex* form.
- We transform them, using new function symbols, to equisatisfiable universal formulas, so called *Skolem variants*.
- We take their *matrices* as axioms of a new theory.

## Conversion rules for quantifiers

Let  $Q$  denote  $\forall$  or  $\exists$  and let  $\bar{Q}$  denote the complementary quantifier.

For every formulas  $\varphi, \psi$  such that  $x$  is not free in the formula  $\psi$ ,

$$\begin{aligned} \models & \quad \neg(Qx)\varphi \leftrightarrow (\bar{Q}x)\neg\varphi \\ \models & \quad ((Qx)\varphi \wedge \psi) \leftrightarrow (Qx)(\varphi \wedge \psi) \\ \models & \quad ((Qx)\varphi \vee \psi) \leftrightarrow (Qx)(\varphi \vee \psi) \\ \models & \quad ((Qx)\varphi \rightarrow \psi) \leftrightarrow (\bar{Q}x)(\varphi \rightarrow \psi) \\ \models & \quad (\psi \rightarrow (Qx)\varphi) \leftrightarrow (Qx)(\psi \rightarrow \varphi) \end{aligned}$$

The above equivalences can be verified semantically or proved by the tableau method (*by taking the universal closure if it is not a sentence*).

**Remark** *The assumption that  $x$  is not free in  $\psi$  is necessary in each rule above (except the first one) for some quantifier  $Q$ . For example,*

$$\not\models ((\exists x)P(x) \wedge P(x)) \leftrightarrow (\exists x)(P(x) \wedge P(x))$$

## Conversion to the prenex normal form

**Proposition** Let  $\varphi'$  be the formula obtained from  $\varphi$  by replacing some occurrences of a subformula  $\psi$  with  $\psi'$ . If  $T \models \psi \leftrightarrow \psi'$ , then  $T \models \varphi \leftrightarrow \varphi'$ .

*Proof* Easily by induction on the structure of the formula  $\varphi$ .  $\square$

**Proposition** For every formula  $\varphi$  there is an equivalent formula  $\varphi'$  in the prenex normal form, i.e.  $\models \varphi \leftrightarrow \varphi'$ .

*Proof* By induction on the structure of  $\varphi$  applying the **conversion rules for quantifiers**, replacing subformulas with their **variants** if needed, and applying the above proposition on equivalent transformations.  $\square$

*For example,*

$$\begin{aligned} ((\forall z)P(x, z) \wedge P(y, z)) &\rightarrow \neg(\exists x)P(x, y) \\ ((\forall u)P(x, u) \wedge P(y, z)) &\rightarrow (\forall x)\neg P(x, y) \\ (\forall u)(P(x, u) \wedge P(y, z)) &\rightarrow (\forall v)\neg P(v, y) \\ (\exists u)((P(x, u) \wedge P(y, z)) &\rightarrow (\forall v)\neg P(v, y)) \\ (\exists u)(\forall v)((P(x, u) \wedge P(y, z)) &\rightarrow \neg P(v, y)) \end{aligned}$$



# Skolem variants

Let  $\varphi$  be a **sentence** of a language  $L$  in the **prenex normal form**, let  $y_1, \dots, y_n$  be the **existentially** quantified variables in  $\varphi$  (in this order), and for every  $i \leq n$  let  $x_1, \dots, x_{n_i}$  be the variables that are **universally** quantified in  $\varphi$  before  $y_i$ . Let  $L'$  be an extension of  $L$  with new  $n_i$ -ary function symbols  $f_i$  for all  $i \leq n$ .

Let  $\varphi_S$  denote the formula of  $L'$  obtained from  $\varphi$  by removing all  $(\exists y_i)$ 's from the prefix and by replacing each occurrence of  $y_i$  with the term  $f_i(x_1, \dots, x_{n_i})$ . Then  $\varphi_S$  is called a **Skolem variant** of  $\varphi$ .

*For example, for the formula  $\varphi$*

$$(\exists y_1)(\forall x_1)(\forall x_2)(\exists y_2)(\forall x_3)R(y_1, x_1, x_2, y_2, x_3)$$

*the following formula  $\varphi_S$  is a Skolem variant of  $\varphi$*

$$(\forall x_1)(\forall x_2)(\forall x_3)R(f_1, x_1, x_2, f_2(x_1, x_2), x_3),$$

*where  $f_1$  is a new constant symbol and  $f_2$  is a new binary function symbol.*

# Properties of Skolem variants

**Lemma** Let  $\varphi$  be a sentence  $(\forall x_1) \dots (\forall x_n)(\exists y)\psi$  of  $L$  and  $\varphi'$  be a sentence  $(\forall x_1) \dots (\forall x_n)\psi(y/f(x_1, \dots, x_n))$  where  $f$  is a new function symbol. Then

- (1) the **reduct**  $\mathcal{A}$  of every model  $\mathcal{A}'$  of  $\varphi'$  to the language  $L$  is a model of  $\varphi$ ,
- (2) every model  $\mathcal{A}$  of  $\varphi$  can be **expanded** into a model  $\mathcal{A}'$  of  $\varphi'$ .

**Remark** Compared to extensions by definition of a function symbol, the expansion in (2) does not need to be unique now.

**Proof** (1) Let  $\mathcal{A}' \models \varphi'$  and  $\mathcal{A}$  be the reduct of  $\mathcal{A}'$  to  $L$ . Since  $\mathcal{A} \models \psi[e(y/a)]$  for every assignment  $e$  where  $a = (f(x_1, \dots, x_n))^{A'}[e]$ , we have also  $\mathcal{A} \models \varphi$ .  
 (2) Let  $\mathcal{A} \models \varphi$ . There exists a function  $f^A: A^n \rightarrow A$  such that for every assignment  $e$  it holds  $\mathcal{A} \models \psi[e(y/a)]$  where  $a = f^A(e(x_1), \dots, e(x_n))$ , and thus the expansion  $\mathcal{A}'$  of  $\mathcal{A}$  by the function  $f^A$  is a model of  $\varphi'$ .  $\square$

**Corollary** If  $\varphi'$  is a Skolem variant of  $\varphi$ , then both statements (1) and (2) hold for  $\varphi, \varphi'$  as well. Hence  $\varphi, \varphi'$  are **equisatisfiable**.

# Skolem's theorem

**Theorem** Every theory  $T$  has an *open conservative extension*  $T^*$ .

*Proof* We may assume that  $T$  is in a closed form. Let  $L$  be its language.

- By replacing each axiom of  $T$  with an equivalent formula in the **prenex normal form** we obtain an equivalent theory  $T^\circ$ .
- By replacing each axiom of  $T^\circ$  with its **Skolem variant** we obtain a theory  $T'$  in an extended language  $L' \supseteq L$ .
- Since the reduct of every model of  $T'$  to the language  $L$  is a model of  $T$ , the theory  $T'$  is an **extension** of  $T$ .
- Furthermore, since every model of  $T$  can be expanded to a model of  $T'$ , it is a **conservative extension**.
- Since every axiom of  $T'$  is a universal sentence, by replacing them with their **matrices** we obtain an open theory  $T^*$  equivalent to  $T'$ .  $\square$

**Corollary** For every theory there is an *equisatisfiable open theory*.

## Reduction of unsatisfiability to propositional logic

If an open theory is unsatisfiable, we can demonstrate it “via ground terms”.

For example, in the language  $L = \langle P, R, f, c \rangle$  the theory

$$T = \{P(x, y) \vee R(x, y), \neg P(c, y), \neg R(x, f(x))\}$$

is unsatisfiable, and this can be demonstrated by an unsatisfiable conjunction of finitely many **instances** of (some) axioms of  $T$  in **ground terms**

$$(P(c, f(c)) \vee R(c, f(c))) \wedge \neg P(c, f(c)) \wedge \neg R(c, f(c)),$$

which may be seen as an unsatisfiable **propositional** formula

$$(p \vee r) \wedge \neg p \wedge \neg r.$$

An instance  $\varphi(x_1/t_1, \dots, x_n/t_n)$  of an open formula  $\varphi$  in free variables  $x_1, \dots, x_n$  is a **ground instance** if all terms  $t_1, \dots, t_n$  are ground terms (i.e. terms without variables).

# Herbrand model

Let  $L = \langle \mathcal{R}, \mathcal{F} \rangle$  be a language with at least one constant symbol. (If needed, we add a new constant symbol to  $L$ .)

- The **Herbrand universe** for  $L$  is the set of all ground terms of  $L$ .  
For example, for  $L = \langle P, f, c \rangle$  with  $f$  binary function sym.,  $c$  constant sym.

$$A = \{c, f(c, c), f(f(c, c), c), f(c, f(c, c)), f(f(c, c), f(c, c)), \dots\}$$

- An  $L$ -structure  $\mathcal{A}$  is a **Herbrand structure** if its domain  $A$  is the Herbrand universe for  $L$  and for each  $n$ -ary function symbol  $f \in \mathcal{F}$ ,  $t_1, \dots, t_n \in A$ ,

$$f^A(t_1, \dots, t_n) = f(t_1, \dots, t_n)$$

(including  $n = 0$ , i.e.  $c^A = c$  for every constant symbol  $c$ ).

**Remark** Compared to a **canonical model**, the relations are not specified.

E.g.  $\mathcal{A} = \langle A, P^A, f^A, c^A \rangle$  with  $P^A = \emptyset$ ,  $c^A = c$ ,  $f^A(c, c) = f(c, c), \dots$

- A **Herbrand model** of a theory  $T$  is a Herbrand structure that models  $T$ .

# Herbrand's theorem

**Theorem** *Let  $T$  be an open theory of a language  $L$  without equality and with at least one constant symbol. Then*

- (a) *either  $T$  has a Herbrand model, or*
- (b) *there are finitely many **ground instances** of axioms of  $T$  whose conjunction is unsatisfiable, and thus  $T$  has no model.*

**Proof** Let  $T'$  be the set of all ground instances of axioms of  $T$ . Consider a finished (e.g. systematic) tableau  $\tau$  from  $T'$  in the language  $L$  (without adding new constant symbols) with the root entry  $F\perp$ .

- If the tableau  $\tau$  contains a noncontradictory branch  $V$ , the canonical model from  $V$  is a Herbrand model of  $T$ .
- Else,  $\tau$  is contradictory, i.e.  $T' \vdash \perp$ . Moreover,  $\tau$  is finite, so  $\perp$  is provable from finitely many formulas of  $T'$ , i.e. their conjunction is unsatisfiable.  $\square$

**Remark** *If the language  $L$  is with equality, we extend  $T$  to  $T^*$  by **axioms of equality** for  $L$  and if  $T^*$  has a Herbrand model  $\mathcal{A}$ , we take its **quotient** by  $=^{\mathcal{A}}$ .*

## Corollaries of Herbrand's theorem

Let  $L$  be a language containing at least one constant symbol.

**Corollary** For every open  $\varphi(x_1, \dots, x_n)$  of  $L$ , the formula  $(\exists x_1) \dots (\exists x_n)\varphi$  is valid if and only if there exist  $mn$  ground terms  $t_{ij}$  of  $L$  for some  $m$  such that

$$\varphi(x_1/t_{11}, \dots, x_n/t_{1n}) \vee \dots \vee \varphi(x_1/t_{m1}, \dots, x_n/t_{mn})$$

is a (propositional) tautology.

**Proof**  $(\exists x_1) \dots (\exists x_n)\varphi$  is valid  $\Leftrightarrow (\forall x_1) \dots (\forall x_n)\neg\varphi$  is unsatisfiable  $\Leftrightarrow \neg\varphi$  is unsatisfiable. The rest follows from Herbrand's theorem for  $\{\neg\varphi\}$ .  $\square$

**Corollary** An open theory  $T$  of  $L$  is satisfiable if and only if the theory  $T'$  of all ground instances of axioms of  $T$  is satisfiable.

**Proof** If  $T$  has a model  $\mathcal{A}$ , every instance of each axiom of  $T$  is valid in  $\mathcal{A}$ , thus  $\mathcal{A}$  is a model of  $T'$ . If  $T$  is unsatisfiable, by H. theorem there are (finitely) formulas of  $T'$  whose conjunction is unsatisfiable, thus  $T'$  is unsatisfiable.  $\square$

# Resolution method in predicate logic - introduction

- A **refutation** procedure - its aim is to show that a given formula (or theory) is unsatisfiable.
- It assumes **open** formulas in **CNF** (and in clausal form).
  - A **literal** is (now) an atomic formula or its negation.
  - A **clause** is a finite set of literals,  $\square$  denotes the **empty clause**.
  - A **formula (in clausal form)** is a (possibly infinite) set of clauses.
- Remark* Every formula (theory) can be converted to an equisatisfiable open formula (theory) in CNF, and then to a formula in clausal form.
- The **resolution rule** is more general - it allows to resolve through literals that are **unifiable**.
- Resolution in predicate logic is based on resolution in **propositional logic** and **unification**.



# Local scope of variables

Variables can be renamed locally within *clauses*.

Let  $\varphi$  be an (*input*) open formula in CNF.

- $\varphi$  is satisfiable if and only if its universal closure  $\varphi'$  is satisfiable.
- For every two formulas  $\psi, \chi$  and a variable  $x$

$$\models (\forall x)(\psi \wedge \chi) \leftrightarrow (\forall x)\psi \wedge (\forall x)\chi$$

(also in the case that  $x$  is free both in  $\psi$  and  $\chi$ ).

- Every clause in  $\varphi$  can thus be replaced by its universal closure.
- We can then take any *variants* of clauses (to rename variables apart).

*For example, by renaming variables in the second clause of (1) we obtain an equisatisfiable formula (2).*

$$(1) \{ \{P(x), Q(x, y)\}, \{\neg P(x), \neg Q(y, x)\} \}$$

$$(2) \{ \{P(x), Q(x, y)\}, \{\neg P(v), \neg Q(u, v)\} \}$$

## Reduction to propositional level (grounding)

*Herbrand's theorem gives us the following (inefficient) method.*

- Let  $S$  be the (input) formula in clausal form.
- We can assume that the language contains at least one constant symbol.
- Let  $S'$  be the set of all **ground instances** of all clauses from  $S$ .
- By introducing propositional letters representing **atomic sentences** we may view  $S'$  as a (possibly infinite) **propositional** formula in clausal form.
- We may verify that it is unsatisfiable by resolution on propositional level.

*For example, for  $S = \{\{P(x, y), R(x, y)\}, \{\neg P(c, y)\}, \{\neg R(x, f(x))\}\}$  the set  $S' = \{\{P(c, c), R(c, c)\}, \{P(c, f(c)), R(c, f(c))\}, \{P(f(c), f(c)), R(f(c), f(c))\}, \dots, \{\neg P(c, c)\}, \{\neg P(c, f(c))\}, \dots, \{\neg R(c, f(c))\}, \{\neg R(f(c), f(f(c)))\}, \dots\}$*

*is unsatisfiable since on propositional level*

$$S' \supseteq \{\{P(c, f(c)), R(c, f(c))\}, \{\neg P(c, f(c))\}, \{\neg R(c, f(c))\}\} \vdash_R \square.$$

# The general resolution rule

Let  $C_1, C_2$  be clauses with **distinct variables** such that

$$C_1 = C'_1 \sqcup \{A_1, \dots, A_n\}, \quad C_2 = C'_2 \sqcup \{\neg B_1, \dots, \neg B_m\},$$

where  $S = \{A_1, \dots, A_n, B_1, \dots, B_m\}$  is unifiable and  $n, m \geq 1$ . Then the clause

$$C = C'_1\sigma \cup C'_2\sigma,$$

where  $\sigma$  is a **most general unification** of  $S$ , is the **resolvent** of  $C_1$  and  $C_2$ .

*For example, in clauses  $\{P(x), Q(x, z)\}$  and  $\{\neg P(y), \neg Q(f(y), y)\}$  we can unify  $S = \{Q(x, z), Q(f(y), y)\}$  applying a most general unification  $\sigma = \{x/f(y), z/y\}$ , and then resolve to a clause  $\{P(f(y)), \neg P(y)\}$ .*

**Remark** *The condition on distinct variables can be satisfied by renaming variables apart. This is sometimes necessary, e.g. from  $\{\{P(x)\}, \{\neg P(f(x))\}\}$  after renaming we can get  $\square$ , but  $\{P(x), P(f(x))\}$  is not unifiable.*

# Resolution proof

We have the same notions as in propositional logic, up to renaming variables.

- **Resolution proof (deduction)** of a clause  $C$  from a formula  $S$  is a **finite** sequence  $C_0, \dots, C_n = C$  such that for every  $i \leq n$ , we have  $C_i = C'_i \sigma$  for some  $C'_i \in S$  and a renaming of variables  $\sigma$ , or  $C_i$  is a resolvent of some previous clauses.
- A clause  $C$  is (resolution) **provable** from  $S$ , denoted by  $S \vdash_R C$ , if it has a resolution proof from  $S$ .
- A (resolution) **refutation** of a formula  $S$  is a resolution proof of  $\square$  from  $S$ .
- $S$  is (resolution) **refutable** if  $S \vdash_R \square$ .

**Remark** Elimination of several literals at once is sometimes necessary, e.g.  $S = \{\{P(x), P(y)\}, \{\neg P(x), \neg P(y)\}\}$  is resolution refutable, but it has no refutation that eliminates only a single literal in each resolution step.

# Resolution in predicate logic - an example

Consider  $T = \{\neg P(x, x), P(x, y) \rightarrow P(y, x), P(x, y) \wedge P(y, z) \rightarrow P(x, z)\}$ .

Is  $T \models (\exists x)\neg P(x, f(x))$ ? Equivalently, is the following  $T'$  unsatisfiable?

$T' = \{\{\neg P(x, x)\}, \{\neg P(x, y), P(y, x)\}, \{\neg P(x, y), \neg P(y, z), P(x, z)\}, \{P(x, f(x))\}\}$

